

$B \rightarrow K^* l^+ l^-$ decay in the two Higgs doublet model with flavor changing neutral currents

T. M. Aliev ,

Physics Department, Middle East Technical University
Ankara, Turkey

E. O. Iltan *

Physics Department, Middle East Technical University
Ankara, Turkey

Abstract

We study the decay width, forward-backward asymmetry and the longitudinal lepton polarization for the exclusive decay $B \rightarrow K^* l^+ l^-$ in the two Higgs doublet model with three level flavor changing neutral currents (model III) and analyse the dependencies of these quantities on the the selected parameters, $\xi^{U,D}$, of model III including the next to leading QCD corrections. It is found that to look for charged Higgs effects, the measurements of the branching ratio, forward-backward asymmetry and the longitudinal lepton polarization for the decay $B \rightarrow K^* l^+ l^-$ are promising.

*E-mail address: eiltan@heraklit.physics.metu.edu.tr

1 Introduction

Rare B meson decays induced by flavor changing neutral current (FCNC) $b \rightarrow s$ transition represent an important channel to test the Standard model (SM) at loop level. They provide a comprehensive information in the determination of the fundamental parameters, such as Cabbibo-Kobayashi-Maskawa (CKM) matrix elements, leptonic decay constants, etc. and open a window to investigate the physics beyond the SM, such as two Higgs Doublet model (2HDM), Minimal Supersymmetric extension of the SM (MSSM) [1], etc. Measurement of the Branching ratios (Br) of the inclusive $B \rightarrow X_s \gamma$ [2] and the exclusive $B \rightarrow K^* \gamma$ [3] decays stimulated the study of the rare B meson decays in a new force.

Currently, the main interest is focused on the rare decays where the SM predicts large Br , which is measurable in the near future. $B \rightarrow K^* l^+ l^-$ decay is one of the candidate of these decays and it is described by $b \rightarrow s l^+ l^-$ transition at the quark level. In the literature [4]- [17] this transition has been investigated extensively in the SM, 2HDM and MSSM. The forward-backward asymmetry of dileptons is another quantity which provides information on the short distance contributions. Further, the longitudinal lepton polarization includes different combinations of the Wilson coefficients C_7^{eff}, C_9^{eff} and C_{10} and it can shed light on the investigation of their signs in the SM and new physics beyond.

The theoretical analysis of exclusive decays is more complicated due to the hadronic form factors, which brings substantial uncertainty in the calculations. However, it is well known that the experimental investigation of inclusive decays is more difficult compared those of exclusive ones. Therefore we will consider the exclusive $B \rightarrow K^* l^+ l^-$ in the present work.

The calculation of the physical observables in the hadronic level needs non-perturbative methods to determine the matrix elements of the quark level effective Hamiltonian between the hadronic states. This problem has been studied in the framework of different approaches such as relativistic quark model by the light-front formalism [17], chiral theory [18], three point QCD sum rules method [19], effective heavy quark theory [20] and light cone QCD sum rules [21, 22].

In this work, we present the next to leading order (NLO) QCD corrected effective Hamiltonian in the 2HDM with flavor changing neutral currents (model III) for the inclusive decay $b \rightarrow s l^+ l^-$ and calculate the differential Br of the exclusive $B \rightarrow K^* l^+ l^-$ decay. Further, we study the forward-backward asymmetry (A_{FB}) and the longitudinal lepton polarization (P_L).

The paper is organized as follows: In Section 2, we give the NLO QCD corrected Hamiltonian and corresponding matrix element for the inclusive $b \rightarrow s \gamma$ decay and calculate the matrix

element. Section 3 is devoted to the calculation of the differential Br , A_{FB} and P_L of the exclusive $B \rightarrow K^* l^+ l^-$ decay. In Section 4 we analyse the dependencies of the Br , A_{FB} and P_L on the Yukawa couplings $\bar{\xi}_{bb}^D$, $\bar{\xi}_{tt}^U$ and estimate the $Br(B \rightarrow K^* l^+ l^-)$ for the model III.

2 QCD corrected short-distance contributions in the model III for the decay $b \rightarrow sl^+ l^-$ with additional long-distance effects

In this section we present the NLO QCD corrections to the inclusive $b \rightarrow sl^+ l^-$ ($l = e, \mu$) decay amplitude in the 2HDM with tree level neutral currents (model III). Before presenting the details of calculations, we would like to give the essential points of this model.

The Yukawa interaction in this general case is

$$\mathcal{L}_Y = \eta_{ij}^U \bar{Q}_{iL} \tilde{\phi}_1 U_{jR} + \eta_{ij}^D \bar{Q}_{iL} \phi_1 D_{jR} + \xi_{ij}^U \bar{Q}_{iL} \tilde{\phi}_2 U_{jR} + \xi_{ij}^D \bar{Q}_{iL} \phi_2 D_{jR} + h.c. , \quad (1)$$

where L and R denote chiral projections $L(R) = 1/2(1 \mp \gamma_5)$, ϕ_i for $i = 1, 2$, are the two scalar doublets, $\eta_{ij}^{U,D}$ and $\xi_{ij}^{U,D}$ are the matrices of the Yukawa couplings. The Flavor Changing (FC) part of the interaction can be written as

$$\mathcal{L}_{Y,FC} = \xi_{ij}^U \bar{Q}_{iL} \tilde{\phi}_2 U_{jR} + \xi_{ij}^D \bar{Q}_{iL} \phi_2 D_{jR} + h.c. , \quad (2)$$

with the choice of ϕ_1 and ϕ_2

$$\phi_1 = \frac{1}{\sqrt{2}} \left[\begin{pmatrix} 0 \\ v + H^0 \end{pmatrix} + \begin{pmatrix} \sqrt{2}\chi^+ \\ i\chi^0 \end{pmatrix} \right] ; \phi_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} \sqrt{2}H^+ \\ H_1 + iH_2 \end{pmatrix} . \quad (3)$$

Here the vacuum expectation values are,

$$\langle \phi_1 \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v \end{pmatrix} ; \langle \phi_2 \rangle = 0 , \quad (4)$$

and the couplings $\xi^{U,D}$ for the FC charged interactions are

$$\begin{aligned} \xi_{ch}^U &= \xi_{neutral} V_{CKM} , \\ \xi_{ch}^D &= V_{CKM} \xi_{neutral} , \end{aligned} \quad (5)$$

where $\xi_{neutral}^{U,D}$ ¹ is defined by the expression

$$\xi_N^{U,D} = (V_L^{U,D})^{-1} \xi^{U,D} V_R^{U,D} . \quad (6)$$

¹In all next discussion we denote $\xi_{neutral}^{U,D}$ as $\xi_N^{U,D}$.

Note that the charged couplings appear as a linear combinations of neutral couplings multiplied by V_{CKM} matrix elements (more details see [23]).

The next step is the calculation of the QCD corrections to $b \rightarrow sl^+l^-$ decay amplitude in the model III. The appropriate framework is that of an effective theory obtained by integrating out the heavy degrees of freedom. In the present case, t quark, $W^\pm$, $H^\pm$, H_1 , and H_2 bosons are the heavy degrees of freedom. Here $H^\pm$ denote charged, H_1 and H_2 denote neutral Higgs bosons. The QCD corrections are done through matching the full theory with the effective low energy theory at the high scale $\mu = m_W$ and evaluating the Wilson coefficients from m_W down to the lower scale $\mu \sim O(m_b)$. Note that we choose the higher scale as $\mu = m_W$ since the evaluation from the scale $\mu = m_{H^\pm}$ to $\mu = m_W$ gives negligible contribution to the Wilson coefficients since the charged Higgs boson is heavy enough from the current theoretical restrictions, for example $m_{H^\pm} \geq 340 \text{ GeV}$ [24], $m_{H^\pm} \geq 480 \text{ GeV}$ [25].

The effective Hamiltonian relevant for our process is

$$\mathcal{H}_{eff} = -4 \frac{G_F}{\sqrt{2}} V_{tb} V_{ts}^* \sum_i C_i(\mu) O_i(\mu), \quad (7)$$

where the O_i are operators given in eqs. (8), (9) and the $C_i(\mu)$ are Wilson coefficients renormalized at the scale μ . The coefficients $C_i(\mu)$ are calculated perturbatively.

The operator basis is the extension of the one used for the SM and 2HDM (model I and II) [26, 27] with the additional flipped chirality partners, similar to the ones used for the $b \rightarrow s\gamma$ decay in the model III [28] and $SU(2)_L \times SU(2)_R \times U(1)$ extensions of the SM [29]:

$$\begin{aligned} O_1 &= (\bar{s}_{L\alpha} \gamma_\mu c_{L\beta}) (\bar{c}_{L\beta} \gamma^\mu b_{L\alpha}), \\ O_2 &= (\bar{s}_{L\alpha} \gamma_\mu c_{L\alpha}) (\bar{c}_{L\beta} \gamma^\mu b_{L\beta}), \\ O_3 &= (\bar{s}_{L\alpha} \gamma_\mu b_{L\alpha}) \sum_{q=u,d,s,c,b} (\bar{q}_{L\beta} \gamma^\mu q_{L\beta}), \\ O_4 &= (\bar{s}_{L\alpha} \gamma_\mu b_{L\beta}) \sum_{q=u,d,s,c,b} (\bar{q}_{L\beta} \gamma^\mu q_{L\alpha}), \\ O_5 &= (\bar{s}_{L\alpha} \gamma_\mu b_{L\alpha}) \sum_{q=u,d,s,c,b} (\bar{q}_{R\beta} \gamma^\mu q_{R\beta}), \\ O_6 &= (\bar{s}_{L\alpha} \gamma_\mu b_{L\beta}) \sum_{q=u,d,s,c,b} (\bar{q}_{R\beta} \gamma^\mu q_{R\alpha}), \\ O_7 &= \frac{e}{16\pi^2} \bar{s}_\alpha \sigma_{\mu\nu} (m_b R + m_s L) b_\alpha \mathcal{F}^{\mu\nu}, \\ O_8 &= \frac{g}{16\pi^2} \bar{s}_\alpha T_{\alpha\beta}^a \sigma_{\mu\nu} (m_b R + m_s L) b_\beta \mathcal{G}^{a\mu\nu}, \\ O_9 &= \frac{e}{16\pi^2} (\bar{s}_{L\alpha} \gamma_\mu b_{L\alpha}) (\bar{l}_\gamma \gamma_\mu l), \\ O_{10} &= \frac{e}{16\pi^2} (\bar{s}_{L\alpha} \gamma_\mu b_{L\alpha}) (\bar{l}_\gamma \gamma_\mu \gamma_5 l), \end{aligned}$$

$$\begin{aligned}
O_{11} &= (\bar{s}_{L\alpha}\gamma_\mu c_{L\beta})(\bar{c}_{R\beta}\gamma^\mu b_{R\alpha}), \\
O_{12} &= (\bar{s}_{L\alpha}\gamma_\mu c_{L\alpha})(\bar{c}_{R\beta}\gamma^\mu b_{R\beta}),
\end{aligned} \tag{8}$$

and the second operator set $O'_1 - O'_{12}$ which are flipped chirality partners of $O_1 - O_{12}$:

$$\begin{aligned}
O'_1 &= (\bar{s}_{R\alpha}\gamma_\mu c_{R\beta})(\bar{c}_{R\beta}\gamma^\mu b_{R\alpha}), \\
O'_2 &= (\bar{s}_{R\alpha}\gamma_\mu c_{R\alpha})(\bar{c}_{R\beta}\gamma^\mu b_{R\beta}), \\
O'_3 &= (\bar{s}_{R\alpha}\gamma_\mu b_{R\alpha}) \sum_{q=u,d,s,c,b} (\bar{q}_{R\beta}\gamma^\mu q_{R\beta}), \\
O'_4 &= (\bar{s}_{R\alpha}\gamma_\mu b_{R\beta}) \sum_{q=u,d,s,c,b} (\bar{q}_{R\beta}\gamma^\mu q_{R\alpha}), \\
O'_5 &= (\bar{s}_{R\alpha}\gamma_\mu b_{R\alpha}) \sum_{q=u,d,s,c,b} (\bar{q}_{L\beta}\gamma^\mu q_{L\beta}), \\
O'_6 &= (\bar{s}_{R\alpha}\gamma_\mu b_{R\beta}) \sum_{q=u,d,s,c,b} (\bar{q}_{L\beta}\gamma^\mu q_{L\alpha}), \\
O'_7 &= \frac{e}{16\pi^2} \bar{s}_\alpha \sigma_{\mu\nu} (m_b L + m_s R) b_\alpha \mathcal{F}^{\mu\nu}, \\
O'_8 &= \frac{g}{16\pi^2} \bar{s}_\alpha T_{\alpha\beta}^a \sigma_{\mu\nu} (m_b L + m_s R) b_\beta \mathcal{G}^{a\mu\nu}, \\
O'_9 &= \frac{e}{16\pi^2} (\bar{s}_{R\alpha}\gamma_\mu b_{R\alpha})(\bar{l}\gamma_\mu l), \\
O'_{10} &= \frac{e}{16\pi^2} (\bar{s}_{R\alpha}\gamma_\mu b_{R\alpha})(\bar{l}\gamma_\mu \gamma_5 l), \\
O'_{11} &= (\bar{s}_{R\alpha}\gamma_\mu c_{R\beta})(\bar{c}_{L\beta}\gamma^\mu b_{L\alpha}), \\
O'_{12} &= (\bar{s}_{R\alpha}\gamma_\mu c_{R\alpha})(\bar{c}_{L\beta}\gamma^\mu b_{L\beta}),
\end{aligned} \tag{9}$$

where α and β are $SU(3)$ colour indices and $\mathcal{F}^{\mu\nu}$ and $\mathcal{G}^{\mu\nu}$ are the field strength tensors of the electromagnetic and strong interactions, respectively.

Denoting the Wilson coefficients for the SM with $C_i^{SM}(m_W)$ and the additional charged Higgs contribution with $C_i^H(m_W)$, we have the initial values for the first set of operators (eq.(8)) [7, 28]

$$\begin{aligned}
C_{1,3,\dots,6,11,12}^{SM}(m_W) &= 0, \\
C_2^{SM}(m_W) &= 1, \\
C_7^{SM}(m_W) &= \frac{3x^3 - 2x^2}{4(x-1)^4} \ln x + \frac{-8x^3 - 5x^2 + 7x}{24(x-1)^3}, \\
C_8^{SM}(m_W) &= -\frac{3x^2}{4(x-1)^4} \ln x + \frac{-x^3 + 5x^2 + 2x}{8(x-1)^3}, \\
C_9^{SM}(m_W) &= -\frac{1}{\sin^2 \theta_W} B(x) + \frac{1 - 4\sin^2 \theta_W}{\sin^2 \theta_W} C(x) - D(x) + \frac{4}{9}, \\
C_{10}^{SM}(m_W) &= \frac{1}{\sin^2 \theta_W} (B(x) - C(x)),
\end{aligned}$$

$$\begin{aligned}
C_{1,\dots,6,11,12}^H(m_W) &= 0, \\
C_7^H(m_W) &= \frac{1}{m_t^2} (\bar{\xi}_{N,tt}^U + \bar{\xi}_{N,tc}^U \frac{V_{cs}^*}{V_{ts}^*}) (\bar{\xi}_{N,tt}^U + \bar{\xi}_{N,tc}^U \frac{V_{cb}}{V_{tb}}) F_1(y), \\
&+ \frac{1}{m_t m_b} (\bar{\xi}_{N,tt}^U + \bar{\xi}_{N,tc}^U \frac{V_{cs}^*}{V_{ts}^*}) (\bar{\xi}_{N,bb}^D + \bar{\xi}_{N,sb}^D \frac{V_{ts}}{V_{tb}}) F_2(y), \\
C_8^H(m_W) &= \frac{1}{m_t^2} (\bar{\xi}_{N,tt}^U + \bar{\xi}_{N,tc}^U \frac{V_{cs}^*}{V_{ts}^*}) (\bar{\xi}_{N,tt}^U + \bar{\xi}_{N,tc}^U \frac{V_{cb}}{V_{tb}}) G_1(y), \\
&+ \frac{1}{m_t m_b} (\bar{\xi}_{N,tt}^U + \bar{\xi}_{N,tc}^U \frac{V_{cs}^*}{V_{ts}^*}) (\bar{\xi}_{N,bb}^D + \bar{\xi}_{N,sb}^D \frac{V_{ts}}{V_{tb}}) G_2(y), \\
C_9^H(m_W) &= \frac{1}{m_t^2} (\bar{\xi}_{N,tt}^U + \bar{\xi}_{N,tc}^U \frac{V_{cs}^*}{V_{ts}^*}) (\bar{\xi}_{N,tt}^U + \bar{\xi}_{N,tc}^U \frac{V_{cb}}{V_{tb}}) H_1(y), \\
C_{10}^H(m_W) &= \frac{1}{m_t^2} (\bar{\xi}_{N,tt}^U + \bar{\xi}_{N,tc}^U \frac{V_{cs}^*}{V_{ts}^*}) (\bar{\xi}_{N,tt}^U + \bar{\xi}_{N,tc}^U \frac{V_{cb}}{V_{tb}}) L_1(y), \tag{10}
\end{aligned}$$

and for the second set of operators eq. (9),

$$\begin{aligned}
C_{1,\dots,12}'^{SM}(m_W) &= 0, \\
C_{1,\dots,6,11,12}'^H(m_W) &= 0, \\
C_7'^H(m_W) &= \frac{1}{m_t^2} (\bar{\xi}_{N,bs}^D \frac{V_{tb}}{V_{ts}^*} + \bar{\xi}_{N,ss}^D) (\bar{\xi}_{N,bb}^D + \bar{\xi}_{N,sb}^D \frac{V_{ts}}{V_{tb}}) F_1(y), \\
&+ \frac{1}{m_t m_b} (\bar{\xi}_{N,bs}^D \frac{V_{tb}}{V_{ts}^*} + \bar{\xi}_{N,ss}^D) (\bar{\xi}_{N,tt}^U + \bar{\xi}_{N,tc}^U \frac{V_{cb}}{V_{tb}}) F_2(y), \\
C_8'^H(m_W) &= \frac{1}{m_t^2} (\bar{\xi}_{N,bs}^D \frac{V_{tb}}{V_{ts}^*} + \bar{\xi}_{N,ss}^D) (\bar{\xi}_{N,bb}^D + \bar{\xi}_{N,sb}^D \frac{V_{ts}}{V_{tb}}) G_1(y), \\
&+ \frac{1}{m_t m_b} (\bar{\xi}_{N,bs}^D \frac{V_{tb}}{V_{ts}^*} + \bar{\xi}_{N,ss}^D) (\bar{\xi}_{N,tt}^U + \bar{\xi}_{N,tc}^U \frac{V_{cb}}{V_{tb}}) G_2(y), \\
C_9'^H(m_W) &= \frac{1}{m_t^2} (\bar{\xi}_{N,bs}^D \frac{V_{tb}}{V_{ts}^*} + \bar{\xi}_{N,ss}^D) (\bar{\xi}_{N,bb}^D + \bar{\xi}_{N,sb}^D \frac{V_{ts}}{V_{tb}}) H_1(y), \\
C_{10}'^H(m_W) &= \frac{1}{m_t^2} (\bar{\xi}_{N,bs}^D \frac{V_{tb}}{V_{ts}^*} + \bar{\xi}_{N,ss}^D) (\bar{\xi}_{N,bb}^D + \bar{\xi}_{N,sb}^D \frac{V_{ts}}{V_{tb}}) L_1(y), \tag{11}
\end{aligned}$$

where $x = m_t^2/m_W^2$ and $y = m_t^2/m_{H^\pm}^2$. In eqs. (10) and (11) we used the redefinition

$$\xi^{U,D} = \sqrt{\frac{4G_F}{\sqrt{2}}} \bar{\xi}^{U,D}. \tag{12}$$

The functions $B(x)$, $C(x)$, $D(x)$, $F_{1(2)}(y)$, $G_{1(2)}(y)$, $H_1(y)$ and $L_1(y)$ are given as

$$\begin{aligned}
B(x) &= \frac{1}{4} \left[\frac{-x}{x-1} + \frac{x}{(x-1)^2} \ln x \right], \\
C(x) &= \frac{x}{4} \left[\frac{x/2-3}{x-1} + \frac{3x/2+1}{(x-1)^2} \ln x \right], \\
D(x) &= \frac{-19x^3/36 + 25x^2/36}{(x-1)^3} + \frac{-x^4/6 + 5x^3/3 - 3x^2 + 16x/9 - 4/9}{(x-1)^4} \ln x,
\end{aligned}$$

$$\begin{aligned}
F_1(y) &= \frac{y(7-5y-8y^2)}{72(y-1)^3} + \frac{y^2(3y-2)}{12(y-1)^4} \ln y , \\
F_2(y) &= \frac{y(5y-3)}{12(y-1)^2} + \frac{y(-3y+2)}{6(y-1)^3} \ln y , \\
G_1(y) &= \frac{y(-y^2+5y+2)}{24(y-1)^3} + \frac{-y^2}{4(y-1)^4} \ln y , \\
G_2(y) &= \frac{y(y-3)}{4(y-1)^2} + \frac{y}{2(y-1)^3} \ln y , \\
H_1(y) &= \frac{1-4\sin^2\theta_W}{\sin^2\theta_W} \frac{xy}{8} \left[\frac{1}{y-1} - \frac{1}{(y-1)^2} \ln y \right] - y \left[\frac{47y^2-79y+38}{108(y-1)^3} - \frac{3y^3-6y+4}{18(y-1)^4} \ln y \right] , \\
L_1(y) &= \frac{1}{\sin^2\theta_W} \frac{xy}{8} \left[-\frac{1}{y-1} + \frac{1}{(y-1)^2} \ln y \right] .
\end{aligned} \tag{13}$$

Note that we neglect the contributions due to the neutral Higgs bosons since their interactions include Yukawa couplings which should be very small (see the discussion section for details).

For the initial values of the Wilson coefficients in the model III (eqs. (10) and (11)), we have

$$\begin{aligned}
C_{1,3,\dots,6,11,12}^{2HDM}(m_W) &= 0 , \\
C_2^{2HDM}(m_W) &= 1 , \\
C_7^{2HDM}(m_W) &= C_7^{SM}(m_W) + C_7^H(m_W) , \\
C_8^{2HDM}(m_W) &= C_8^{SM}(m_W) + C_8^H(m_W) , \\
C_9^{2HDM}(m_W) &= C_9^{SM}(m_W) + C_9^H(m_W) , \\
C_{10}^{2HDM}(m_W) &= C_{10}^{SM}(m_W) + C_{10}^H(m_W) , \\
\\
C'_{1,2,3,\dots,6,11,12}{}^{2HDM}(m_W) &= 0 , \\
C'_7{}^{2HDM}(m_W) &= C'_7{}^{SM}(m_W) + C'_7{}^H(m_W) , \\
C'_8{}^{2HDM}(m_W) &= C'_8{}^{SM}(m_W) + C'_8{}^H(m_W) , \\
C'_9{}^{2HDM}(m_W) &= C'_9{}^{SM}(m_W) + C'_9{}^H(m_W) , \\
C'_{10}{}^{2HDM}(m_W) &= C'_{10}{}^{SM}(m_W) + C'_{10}{}^H(m_W) .
\end{aligned} \tag{14}$$

Using the initial values, we can calculate the coefficients C_i^{2HDM} and $C_i'^{2HDM}$ at any lower scale with five quark effective theory where large logarithms can be summed using the renormalization group and their evaluations are similar to the SM case.

The operators O_5, O_6, O_{11} and O_{12} (O'_5, O'_6, O'_{11} and O'_{12}) give a contribution to the leading order matrix element of $b \rightarrow s\gamma$ and the magnetic moment type coefficient $C_7^{eff}(\mu)$ ($C_7'^{eff}(\mu)$)

is redefined in the NDR scheme as [28]:

$$\begin{aligned}
C_7^{eff}(\mu) &= C_7^{2HDM}(\mu) + Q_d (C_5^{2HDM}(\mu) + N_c C_6^{2HDM}(\mu)) , \\
&+ Q_u \left(\frac{m_c}{m_b} C_{12}^{2HDM}(\mu) + N_c \frac{m_c}{m_b} C_{11}^{2HDM}(\mu) \right) , \\
C_7'^{eff}(\mu) &= C_7'^{2HDM}(\mu) + Q_d (C_5'^{2HDM}(\mu) + N_c C_6'^{2HDM}(\mu)) \\
&+ Q_u \left(\frac{m_c}{m_b} C_{12}'^{2HDM}(\mu) + N_c \frac{m_c}{m_b} C_{11}'^{2HDM}(\mu) \right) .
\end{aligned} \tag{15}$$

The NLO corrected coefficients $C_7^{2HDM}(\mu)$ and $C_7'^{2HDM}(\mu)$ are given as

$$\begin{aligned}
C_7^{2HDM}(\mu) &= C_7^{LO,2HDM}(\mu) + \frac{\alpha_s(\mu)}{4\pi} C_7^{(1)2HDM}(\mu) , \\
C_7'^{2HDM}(\mu) &= C_7'^{LO,2HDM}(\mu) + \frac{\alpha_s(\mu)}{4\pi} C_7'^{(1)2HDM}(\mu) .
\end{aligned} \tag{16}$$

The functions $C_7^{LO,2HDM}(\mu)$ and $C_7'^{LO,2HDM}(\mu)$ are the leading order QCD corrected Wilson coefficients [15, 26, 27]:

$$\begin{aligned}
C_7^{LO,2HDM}(\mu) &= \eta^{16/23} C_7^{2HDM}(m_W) + (8/3)(\eta^{14/23} - \eta^{16/23}) C_8^{2HDM}(m_W) \\
&+ C_2^{2HDM}(m_W) \sum_{i=1}^8 h_i \eta^{a_i} , \\
C_7'^{LO,2HDM}(\mu) &= \eta^{16/23} C_7'^{2HDM}(m_W) + (8/3)(\eta^{14/23} - \eta^{16/23}) C_8'^{2HDM}(m_W)
\end{aligned} \tag{17}$$

where $\eta = \alpha_s(m_W)/\alpha_s(\mu)$, h_i and a_i are the numbers which appear during the evaluation [15]. $C_7^{(1)2HDM}(\mu)$ is the α_s correction to the leading order result that its explicit form can be found in [24, 30]. $C_7'^{(1)2HDM}(\mu)$ can be obtained by replacing the Wilson coefficients in $C_7^{(1)2HDM}(\mu)$ with their primed counterparts.

Since O_2 (O_2') produce dilepton via virtual photon, its Wilson coefficient $C_2(\mu)$ ($C_2'(\mu)$) and the coefficients $C_1(\mu)$, $C_3(\mu)$, ..., $C_6(\mu)$ ($C_2'(\mu)$, $C_3'(\mu)$, ..., $C_6'(\mu)$) induced by the operator mixing, give contributions to $C_9^{eff}(\mu)$ ($C_9'^{eff}(\mu)$). The perturbative part of $C_9^{eff}(\mu)$ [15, 27] and $C_9'^{eff}(\mu)$ including NLO QCD corrections are defined in the NDR scheme as:

$$\begin{aligned}
C_9^{pert}(\mu) &= C_9^{2HDM}(\mu) \tilde{\eta}(\hat{s}) \\
&+ h(z, \hat{s}) (3C_1(\mu) + C_2(\mu) + 3C_3(\mu) + C_4(\mu) + 3C_5(\mu) + C_6(\mu)) \\
&- \frac{1}{2} h(1, \hat{s}) (4C_3(\mu) + 4C_4(\mu) + 3C_5(\mu) + C_6(\mu)) \\
&- \frac{1}{2} h(0, \hat{s}) (C_3(\mu) + 3C_4(\mu)) + \frac{2}{9} (3C_3(\mu) + C_4(\mu) + 3C_5(\mu) + C_6(\mu)) ,
\end{aligned} \tag{18}$$

and

$$\begin{aligned}
C_9'^{pert}(\mu) &= C_9'^{2HDM}(\mu) \tilde{\eta}(\hat{s}) \\
&+ h(z, \hat{s}) (3C_1'(\mu) + C_2'(\mu) + 3C_3'(\mu) + C_4'(\mu) + 3C_5'(\mu) + C_6'(\mu)) \\
&- \frac{1}{2} h(1, \hat{s}) (4C_3'(\mu) + 4C_4'(\mu) + 3C_5'(\mu) + C_6'(\mu)) \\
&- \frac{1}{2} h(0, \hat{s}) (C_3'(\mu) + 3C_4'(\mu)) + \frac{2}{9} (3C_3'(\mu) + C_4'(\mu) + 3C_5'(\mu) + C_6'(\mu)) .
\end{aligned} \tag{19}$$

where $z = \frac{m_c}{m_b}$ and $\hat{s} = \frac{q^2}{m_b^2}$. In the above expression $\tilde{\eta}(\hat{s})$ represents the one gluon correction to the matrix element O_9 with $m_s = 0$ [27] and the function $h(z, \hat{s})$ arises from the one loop contributions of the four quark operators $O_1, \dots, O_6$ ($O_1', \dots, O_6'$). Their explicit forms can be found in [15, 27]. In addition to the short distance part there exist the long distance (LD) effects due to the real $c\bar{c}$ in the intermediate states, i.e. the cascade process $B \rightarrow K^* \psi_i \rightarrow K^* l^+ l^-$ where $i = 1, \dots, 6$. In the current literature, there are four different approaches to take these intermediate states into account: HQET approach [31], AMM approach [12], LSW approach [32] and KS approach [33]. In our calculations, we follow the approach AMM. (We also use KS approach in our numerical analysis and see that results obtained in both methods are closed each other.) In this method the resonance $c\bar{c}$ contribution is parametrized using a Breit-Wigner shape with normalizations fixed by data, and this contribution is added to the perturbative one coming from the $c\bar{c}$ loop:

$$C_9^{eff}(\mu) = C_9^{pert}(\mu) + Y_{reson}(\hat{s}) , \tag{20}$$

where $Y_{reson}(\hat{s})$ in NDR scheme is defined as

$$\begin{aligned}
Y_{reson}(\hat{s}) &= -\frac{3}{\alpha_{em}^2} \kappa \sum_{V_i=\psi_i} \frac{\pi \Gamma(V_i \rightarrow ll) m_{V_i}}{q^2 - m_{V_i}^2 + im_{V_i} \Gamma_{V_i}} \\
&(3C_1(\mu) + C_2(\mu) + 3C_3(\mu) + C_4(\mu) + 3C_5(\mu) + C_6(\mu)) .
\end{aligned} \tag{21}$$

For the expression $C_9'^{eff}(\mu)$, it is enough to replace all unprimed coefficients with primed ones. In eqs. (18) and (19) the phenomenological parameter $\kappa = 2.3$ is chosen to be able to reproduce the correct value of the branching ratio $Br(B \rightarrow J/\psi X \rightarrow X l \bar{l}) = Br(B \rightarrow J/\psi X) Br(J/\psi \rightarrow X l \bar{l})$ [12]. The NLO corrected coefficients $C_i, i = 1, \dots, 6$ can be found in [24, 30]. In our numerical calculations we neglect the coefficients $C_5^{2HDM}(\mu)$, $C_6^{2HDM}(\mu)$ and $C_5'^{2HDM}(\mu)$, $C_6'^{2HDM}(\mu)$ since they are numerically small at m_b scale.

Note that the expressions for unprimed Wilson coefficients in our case can be obtained from the results in [30] by the following replacements:

$$|Y|^2 \rightarrow \frac{1}{m_t^2} (\bar{\xi}_{N,tt}^U + \bar{\xi}_{N,tc}^U \frac{V_{cs}^*}{V_{ts}^*}) (\bar{\xi}_{N,tt}^U + \bar{\xi}_{N,tc}^U \frac{V_{cb}}{V_{tb}}) ,$$

$$XY \rightarrow \frac{1}{m_t m_b} (\bar{\xi}_{N,tt}^U + \bar{\xi}_{N,tc}^U \frac{V_{cs}^*}{V_{ts}^*}) (\bar{\xi}_{N,bb}^D + \bar{\xi}_{N,sb}^D \frac{V_{ts}}{V_{tb}}) \quad (22)$$

To obtain primed coefficients, it is enough to use the primed ones at m_W level (eq. (11)) since the evaluation of $C'_i(\mu)$ from $\mu = m_W$ to $\mu = m_b$ is the same as that of $C_i(\mu)$.

For model II (model I) Y and XY are

$$\begin{aligned} Y &= 1/\tan\beta \ (1/\tan\beta) , \\ XY &= 1 \ (-1/\tan^2\beta) . \end{aligned} \quad (23)$$

At this stage we would like to make the following remark. In [34] QED corrections to the Wilson coefficient C_7^{eff} for $B \rightarrow X_s \gamma$ decay are calculated in addition to the NLO QCD ones and it was shown that QED corrections were almost one order smaller than NLO QCD ones. We expect that the similar situation exists for C_9^{eff} . Therefore, in our calculations we neglect them.

Finally, neglecting the strange quark mass, the matrix element for $b \rightarrow s \ell^+ \ell^-$ decay is obtained as:

$$\begin{aligned} \mathcal{M} &= -\frac{G_F \alpha_{em}}{2\sqrt{2}\pi} V_{tb} V_{ts}^* \left\{ \left(C_9^{eff}(\mu) \bar{s} \gamma_\mu (1 - \gamma_5) b + C_9'^{eff}(\mu) \bar{s} \gamma_\mu (1 + \gamma_5) b \right) \bar{\ell} \gamma^\mu \ell \right. \\ &+ \left(C_{10}(\mu) \bar{s} \gamma_\mu (1 - \gamma_5) b + C_{10}'(\mu) \bar{s} \gamma_\mu (1 + \gamma_5) b \right) \bar{\ell} \gamma^\mu \gamma_5 \ell \\ &\left. - 2 \left(C_7^{eff}(\mu) \frac{m_b}{q^2} \bar{s} i \sigma_{\mu\nu} q^\nu (1 + \gamma_5) b + C_7'^{eff}(\mu) \frac{m_b}{q^2} \bar{s} i \sigma_{\mu\nu} q^\nu (1 - \gamma_5) b \right) \bar{\ell} \gamma^\mu \ell \right\} . \end{aligned} \quad (24)$$

3 The exclusive $B \rightarrow K^* l^+ l^-$ decay

Now, our aim is to look at the problem from the hadronic side. To calculate the decay width, branching ratio, etc., for the exclusive $B \rightarrow K^* l^+ l^-$ decay, we need the matrix elements $\langle K^* | \bar{s} \gamma_\mu (1 \pm \gamma_5) b | B \rangle$, and $\langle K^* | \bar{s} i \sigma_{\mu\nu} q^\nu (1 \pm \gamma_5) b | B \rangle$. Using the parametrization of the form factors as in [19], the matrix element of the $B \rightarrow K^* l^+ l^-$ decay is obtained as [21]:

$$\begin{aligned} \mathcal{M} &= -\frac{G \alpha_{em}}{2\sqrt{2}\pi} V_{tb} V_{ts}^* \left\{ \bar{\ell} \gamma^\mu \ell \left[2A_{tot} \epsilon_{\mu\nu\rho\sigma} \epsilon^{*\nu} p_{K^*}^\rho q^\sigma + iB_{1tot} \epsilon_\mu^* - iB_{2tot}(\epsilon^* q)(p_B + p_{K^*})_\mu - iB_{3tot}(\epsilon^* q) q_\mu \right] \right. \\ &+ \left. \bar{\ell} \gamma^\mu \gamma_5 \ell \left[2C_{tot} \epsilon_{\mu\nu\rho\sigma} \epsilon^{*\nu} p_{K^*}^\rho q^\sigma + iD_{1tot} \epsilon_\mu^* - iD_{2tot}(\epsilon^* q)(p_B + p_{K^*})_\mu - iD_{3tot}(\epsilon^* q) q_\mu \right] \right\} , \end{aligned} \quad (25)$$

where $\epsilon^{*\mu}$ is the polarization vector of K^* meson, p_B and p_{K^*} are four momentum vectors of B and K^* mesons, $q = p_B - p_{K^*}$ and

$$A_{tot} = A + A' ,$$

$$\begin{aligned}
B_{1tot} &= B_1 + B'_1 , \\
B_{2tot} &= B_2 + B'_2 , \\
B_{3tot} &= B_3 + B'_3 , \\
C_{tot} &= C + C' , \\
D_{1tot} &= D_1 + D'_1 , \\
D_{2tot} &= D_2 + D'_2 , \\
D_{3tot} &= D_3 + D'_3 .
\end{aligned} \tag{26}$$

Here

$$\begin{aligned}
A &= -C_9^{eff} \frac{V}{m_B + m_{K^*}} - 4C_7^{eff} \frac{m_b}{q^2} T_1 , \\
B_1 &= -C_9^{eff} (m_B + m_{K^*}) A_1 - 4C_7^{eff} \frac{m_b}{q^2} (m_B^2 - m_{K^*}^2) T_2 , \\
B_2 &= -C_9^{eff} \frac{A_2}{m_B + m_{K^*}} - 4C_7^{eff} \frac{m_b}{q^2} \left(T_2 + \frac{q^2}{m_B^2 - m_{K^*}^2} T_3 \right) , \\
B_3 &= -C_9^{eff} \frac{2m_{K^*}}{q^2} (A_3 - A_0) + 4C_7^{eff} \frac{m_b}{q^2} T_3 , \\
C &= -C_{10} \frac{V}{m_B + m_{K^*}} , \\
D_1 &= -C_{10} (m_B + m_{K^*}) A_1 , \\
D_2 &= -C_{10} \frac{A_2}{m_B + m_{K^*}} , \\
D_3 &= -C_{10} \frac{2m_{K^*}}{q^2} (A_3 - A_0) ,
\end{aligned} \tag{27}$$

and

$$\begin{aligned}
A' &= -C_9'^{eff} \frac{V}{m_B + m_{K^*}} - 4C_7'^{eff} \frac{m_b}{q^2} T_1 , \\
B'_1 &= C_9'^{eff} (m_B + m_{K^*}) A_1 + 4C_7'^{eff} \frac{m_b}{q^2} (m_B^2 - m_{K^*}^2) T_2 , \\
B'_2 &= C_9'^{eff} \frac{A_2}{m_B + m_{K^*}} + 4C_7'^{eff} \frac{m_b}{q^2} \left(T_2 + \frac{q^2}{m_B^2 - m_{K^*}^2} T_3 \right) , \\
B'_3 &= C_9'^{eff} \frac{2m_{K^*}}{q^2} (A_3 - A_0) - 4C_7'^{eff} \frac{m_b}{q^2} T_3 , \\
C' &= -C_{10}' \frac{V}{m_B + m_{K^*}} , \\
D'_1 &= C_{10}' (m_B + m_{K^*}) A_1 , \\
D'_2 &= C_{10}' \frac{A_2}{m_B + m_{K^*}} ,
\end{aligned}$$

$$D'_3 = C'_{10} \frac{2m_{K^*}}{q^2} (A_3 - A_0) , \quad (28)$$

The hadronic formfactors V , A_1 , A_2 , A_0 , T_1 , T_2 and T_3 has been calculated in the framework of light-cone QCD sum rules in [21, 22]. In our calculations we use the results of [22] where radiative corrections to the leading twist wave function and $SU(3)$ breaking effects are also taken into account. The q^2 dependence of the form factors can be represented in terms of three parameters as:

$$F(q^2) = \frac{F(0)}{1 - a_F \frac{q^2}{m_B^2} + b_F \left(\frac{q^2}{m_B^2}\right)^2} , \quad (29)$$

where the values of parameters $F(0)$, a_F and b_F are listed in Table 2.

	$F(0)$	a_F	b_F
A_1	0.34 ± 0.05	0.60	-0.023
A_2	0.28 ± 0.04	1.18	0.281
V	0.46 ± 0.07	1.55	0.575
T_1	0.19 ± 0.03	1.59	0.615
T_2	0.19 ± 0.03	0.49	-0.241
T_3	0.13 ± 0.02	1.20	0.098

Table 1: The values of parameters existing in eq.(29) for the various form factors of the transition $B \rightarrow K^*$.

Light cone QCD sum rules is applicable in the region, $m_b^2 - q^2 \sim \text{few } GeV^2$ and this leads to $q_{max}^2 \sim 20 GeV^2$. To extend the results to the full physical region, we use the above parametrization in such a way that it correctly reproduces light cone QCD sum rules results up to $q_{max}^2 \sim 20 GeV^2$. All the errors come from uncertainty of b-quark mass, higher twist wave functions, and variation of Borel parameter are added quadratically.

Using eq.(25) we get the double differential decay rate:

$$\begin{aligned} \frac{d\Gamma}{dq^2 dz} = & \frac{G^2 \alpha_{em}^2 |V_{tb} V_{ts}^*|^2 \lambda^{1/2}}{2^{12} \pi^5 m_B} \left\{ 2\lambda m_B^4 \left[m_B^2 s (1 + z^2) (|A_{tot}|^2 + |C_{tot}|^2) \right] \right. \\ & + \frac{\lambda m_B^4}{2r} \left[\lambda m_B^2 (1 - z^2) (|B_{2tot}|^2 + |D_{2tot}|^2) \right] \\ & + \frac{1}{2r} \left[m_B^2 \{ \lambda (1 - z^2) + 8rs \} (|B_{1tot}|^2 + |D_{1tot}|^2) \right. \\ & - 2\lambda m_B^4 (1 - r - s) (1 - z^2) \{ \text{Re}(B_{1tot} B_{2tot}^*) + \text{Re}(D_{1tot} D_{2tot}^*) \} \left. \right] \\ & - 8m_B^4 s \lambda^{1/2} z \left[\{ \text{Re}(B_{1tot} C_{tot}^*) + \text{Re}(A_{tot} D_{1tot}^*) \} \right] \left. \right\} , \quad (30) \end{aligned}$$

where $z = \cos\theta$, θ is the angle between the momentum of ℓ lepton and that of B meson in the center of mass frame of the lepton pair, $\lambda = 1 + r^2 + s^2 - 2r - 2s - 2rs$, $r = \frac{m_{K^*}^2}{m_B^2}$ and $s = \frac{q^2}{m_B^2}$.

As we noted, A_{FB} and P_L can give more precise information about the Wilson coefficients C_7^{eff} , C_9^{eff} and C_{10} . They are defined as:

$$A_{FB}(q^2) = \frac{\int_0^1 dz \frac{d\Gamma}{dq^2 dz} - \int_{-1}^0 dz \frac{d\Gamma}{dq^2 dz}}{\int_0^1 dz \frac{d\Gamma}{dq^2 dz} + \int_{-1}^0 dz \frac{d\Gamma}{dq^2 dz}} \quad (31)$$

and

$$P_L(q^2) = \frac{\frac{d\Gamma}{dq^2}(\xi = -1) - \frac{d\Gamma}{dq^2}(\xi = +1)}{\frac{d\Gamma}{dq^2}(\xi = -1) + \frac{d\Gamma}{dq^2}(\xi = +1)} \quad (32)$$

where $\xi = -1(+1)$ corresponds to the left (right) handed lepton in the final state. After the standard calculation, we get

$$P_L = \frac{1}{\Delta} \left\{ \frac{32}{3} s \lambda m_B^6 \text{Re}(A_{tot} C_{tot}^*) + \frac{4}{3} \frac{\lambda^2 m_B^6}{r} \text{Re}(B_{2tot} D_{2tot}^*) + 4 \frac{m_B^2}{r} \left[\frac{\lambda}{3} + 4rs \right] \text{Re}(B_{1tot} D_{1tot}^*) \right. \\ \left. - \frac{4}{3} \frac{\lambda m_B^4}{r} (1 - r - s) [\text{Re}(B_{2tot} D_{1tot}^*) + \text{Re}(B_{1tot} D_{2tot}^*)] \right\}. \quad (33)$$

The denominator Δ in eq. (33) can be obtained from eq.(30) by integration over z of the terms within the curly bracket. In our numerical analysis we used the input values given in Table (2).

Parameter	Value
m_c	1.4 (GeV)
m_b	4.8 (GeV)
α_{em}^{-1}	129
λ_t	0.04
$\Gamma_{tot}(B_d)$	$3.96 \cdot 10^{-13}$ (GeV)
m_{B_d}	5.28 (GeV)
m_t	175 (GeV)
m_W	80.26 (GeV)
m_Z	91.19 (GeV)
Λ_{QCD}	0.214 (GeV)
$\alpha_s(m_Z)$	0.117
$\sin\theta_W$	0.2325

Table 2: The values of the input parameters used in the numerical calculations.

4 Discussion

Before we present our numerical results, we would like to discuss briefly the free parameters of the model III. This model induces many free parameters, such as $\xi_{ij}^{U,D}$ where i, j are flavor indices and it is necessary to restrict them using the experimental measurements. The contributions of the neutral Higgs bosons h_0 and A_0 to the Wilson coefficient C_7 (see the appendix of [37] for details) are

$$\begin{aligned} C_7^{h_0}(m_W) &= (V_{tb}V_{ts}^*)^{-1} \sum_{i=d,s,b} \bar{\xi}_{N,bi}^D \bar{\xi}_{N,is}^D \frac{Q_i}{8 m_i m_b}, \\ C_7^{A_0}(m_W) &= (V_{tb}V_{ts}^*)^{-1} \sum_{i=d,s,b} \bar{\xi}_{N,bi}^D \bar{\xi}_{N,is}^D \frac{Q_i}{8 m_i m_b}, \end{aligned} \quad (34)$$

where m_i and Q_i are the masses and charges of the down quarks ($i = d, s, b$) respectively. These expressions show that the neutral Higgs bosons can give a large contribution to the coefficient C_7 which is in contradiction with the CLEO data announced recently [38],

$$Br(B \rightarrow X_s \gamma) = (3.15 \pm 0.35 \pm 0.32) 10^{-4}. \quad (35)$$

Such dangerous terms are removed with the assumption that the couplings $\bar{\xi}_{N,is}^D (i = d, s, b)$ and $\bar{\xi}_{N,db}^D$ are negligible to be able to reach the conditions $\bar{\xi}_{N,bb}^D \bar{\xi}_{N,is}^D \ll 1$ and $\bar{\xi}_{N,db}^D \bar{\xi}_{N,ds}^D \ll 1$. With the discussion given above and using the constraints [28], coming from the $\Delta F = 2$ mixing, the ρ parameter [39], and the measurement by CLEO Collaboration we get following restrictions: $\bar{\xi}_{Ntc}^D \ll \bar{\xi}_{Ntt}^U$, $\bar{\xi}_{Nbb}^D$ and $\bar{\xi}_{Nib}^D \sim 0$, $\bar{\xi}_{Nij}^D \sim 0$, where the indices i, j denote d and s quarks. Under this assumption, the Wilson coefficients C_7' , C_9' and C_{10}' can be neglected compared to unprimed ones and the neutral Higgs contributions are suppressed because the Yukawa vertices are the combinations of $\bar{\xi}_{Nsb}^D$ and $\bar{\xi}_{Nss}^D$. After these preliminary remarks, let us start with our numerical analysis.

In this section, we study the q^2 dependencies of the differential Br , A_{FB} and P_L of the decay $B \rightarrow K^* l^+ l^-$ for the selected parameters of the model III ($\bar{\xi}_{Ntt}^U$, $\bar{\xi}_{Nbb}^D$). In figs. 1 and 2 we plot the differential Br of the decay $B \rightarrow K^* l^+ l^-$ with respect to the dilepton mass q^2 for $\bar{\xi}_{N,bb}^D = 40 m_b$ and charged Higgs mass $m_{H^\pm} = 400 GeV$ at the scale $\mu = m_b$. Fig. 1 represents the case where the ratio $|r_{tb}| = |\frac{\bar{\xi}_{N,tt}^U}{\bar{\xi}_{N,bb}^D}| \ll 1$. In this case, it is shown that the differential Br obtained in the model III almost coincides with the one calculated in the SM. In Fig. 2, we present the same dependence as in Fig. 1 for the case $r_{tb} \gg 1$ and $\bar{\xi}_{N,bb}^D = 40 m_b$. Here $\bar{\xi}_{N,tt}^U$ lies in the restriction region coming from the constraints mentioned above. The strong enhancement is observed compared to the SM result especially for the small values of q^2 . Further, the differential Br increases with the increasing $\bar{\xi}_{N,bb}^D$ for $r_{tb} \gg 1$.

Figs. 3 and 4 show the q^2 dependence of A_{FB} for $\bar{\xi}_{N,bb}^D = 40 m_b$ and charged Higgs mass $m_{H^\pm} = 400 \text{ GeV}$ at the scale $\mu = m_b$. For $|r_{tb}| \ll 1$ (Fig. 3) A_{FB} changes its sign, however for $r_{tb} \gg 1$ (Fig. 4) it is positive without LD effects. Therefore the determination of the sign of A_{FB} in the region $0 \leq q^2 \leq 2.8 (\text{GeV})^2$ (here the upper limit corresponds to the value where A_{FB} change sign in the SM) can give a unique information about the existence of the model III. Further, A_{FB} becomes more positive with increasing $\bar{\xi}_{N,bb}^D$. It is interesting to note that the final lepton tends to move almost in the direction of B meson with increasing $\bar{\xi}_{N,bb}^D$ for $r_{tb} \gg 1$, especially in the q^2 region far from LD effects.

The behaviour of P_L versus q^2 is presented in series of figures (5 and 6) for $\bar{\xi}_{N,bb}^D$ and charged Higgs mass $m_{H^\pm} = 400 \text{ GeV}$ at the scale $\mu = m_b$. P_L becomes more negative in the region $r_{tb} \gg 1$ (Fig. 6) compared to the one in the region $|r_{tb}| \ll 1$ (Fig. 5). This tendency increases with increasing $\bar{\xi}_{N,bb}^D$.

Now we would like to present the values of Br for the $B \rightarrow K^* l^+ l^-$ decay in the SM and model III, without the LD effects. After integrating over q^2 , we get

$$Br(B \rightarrow K^* l^+ l^-) = 0.278 \pm 0.050 \times 10^{-5} \text{ (SM)} \quad (36)$$

and for the model III

$$Br(B \rightarrow K^* l^+ l^-) = \begin{cases} 0.278 \pm 0.050 \times 10^{-5} & (|r_{tb}| \ll 1) \\ 0.990 \pm 0.050 \times 10^{-5} & (r_{tb} \gg 1, \bar{\xi}_{N,bb}^D = 40 m_b). \end{cases} \quad (37)$$

The Br in the SM is almost equal to the model III value for $|r_{tb}| \ll 1$. However, the strong enhancement can be observed for $r_{tb} \gg 1$, especially with increasing $\bar{\xi}_{N,bb}^D$. Using the CLEO data for the inclusive branching ratio $Br(B \rightarrow X_s l^+ l^-) \leq 0.42 \times 10^{-4}$ [40], it is possible to estimate an upper limit for $\bar{\xi}_{N,bb}^D$, which is $\bar{\xi}_{N,bb}^D \leq 40 m_b$.

In conclusion, we analyse the selected model III parameters ($\bar{\xi}_{N,bb}^D, \bar{\xi}_{N,tt}^U$) dependencies of the differential Br , A_{FB} and P_L of the decay $B \rightarrow K^* l^+ l^-$. We obtain that the strong enhancement of the differential Br is possible in the framework of the model III and observe that A_{FB} and P_L are very sensitive to the model III parameters ($\bar{\xi}_{N,bb}^D, \bar{\xi}_{N,tt}^U$). Therefore, their experimental investigations ensure a crucial test for new physics.

References

- [1] J. L. Hewett, in Proc. of the 21st Annual SLAC Summer Institute, ed. L. De Porcel and C. Dunwoode, SLAC-PUB-6521 (1994)
- [2] M. S. Alam *et. al.* **CLEO** Collaboration, *Phys. Rev.Lett.* **74** (1995) 2885;
- [3] R. Ammar *et. al.* **CLEO** Collaboration, *Phys. Rev. Lett.* **71** (1993) 674.
- [4] W. -S. Hou, R. S. Willey and A. Soni, *Phys. Rev. Lett.* **58** (1987) 1608.
- [5] N. G. Deshpande and J. Trampetic, *Phys. Rev. Lett.* **60** (1988) 2583.
- [6] C. S. Lim, T. Morozumi and A. I. Sanda, *Phys. Lett.* **B218** (1989) 343.
- [7] B. Grinstein, M. J. Savage and M. B. Wise, *Nucl. Phys.* **B319** (1989) 271.
- [8] C. Dominguez, N. Paver and Riazuddin, *Phys. Lett.* **B214** (1988) 459.
- [9] N. G. Deshpande, J. Trampetic and K. Ponose, *Phys. Rev.* **D39** (1989) 1461.
- [10] P. J. O'Donnell and H. K. Tung, *Phys. Rev.* **D43** (1991) 2067.
- [11] N. Paver and Riazuddin, *Phys. Rev.* **D45** (1992) 978.
- [12] A. Ali, T. Mannel and T. Morozumi, *Phys. Lett.* **B273** (1991) 505.
- [13] A. Ali, G. F. Giudice and T. Mannel, *Z. Phys.* **C67** (1995) 417.
- [14] C. Greub, A. Ioannissian and D. Wyler, *Phys. Lett.* **B346** (1995) 145;
D. Liu *Phys. Lett.* **B346** (1995) 355;
G. Burdman, *Phys. Rev.* **D52** (1995) 6400;
Y. Okada, Y. Shimizu and M. Tanaka *Phys. Lett.* **B405** (1997) 297.
- [15] A. J. Buras and M. Münz, *Phys. Rev.* **D52** (1995) 186.
- [16] N. G. Deshpande, X. -G. He and J. Trampetic, *Phys. Lett.* **B367** (1996) 362.
- [17] W. Jaus and D. Wyler, *Phys. Rev.* **D41** (1990) 3405.
- [18] R. Casalbuoni, A. Deandra, N. Di Bartolemo, R. Gatto and G. Nardulli,
Phys. Lett. **B312** (1993) 315.

- [19] P. Colangelo, F. De Fazio, P. Santorelli and E. Scrimieri, *Phys. Rev.* **D53** (1996) 3672.
- [20] W. Roberts, *Phys. Rev.* **D54** (1996) 863.
- [21] T. M. Aliev, A. Özpineci and M. Savcı, *Phys. Rev.* **D56** (1997) 4260.
- [22] P. Ball and V. Braun, *Phys. Rev.* **D57** (1998) 4260.
- [23] D. Atwood, L. Reina and A. Soni, *Phys. Rev.* **D53** (1996) 119.
- [24] M. Ciuchini, G. Degrossi, P. Gambino and G. I. Giudice, *Nucl. Phys.* **B527** (1998) 21.
- [25] T. M. Aliev, G. Hiller and E. O. Iltan, *Nucl. Phys.* **B515** (1998) 321.
- [26] B. Grinstein, R. Springer, and M. Wise, *Nucl. Phys.* **B339** (1990) 269; R. Grigjanis, P.J. O'Donnel, M. Sutherland and H. Navelet, *Phys. Lett.* **B213** (1988) 355; *Phys. Lett.* **B286** (1992) E, 413; G. Cella, G. Curci, G. Ricciardi and A. Viceré, *Phys. Lett.* **B325** (1994) 227, *Nucl. Phys.* **B431** (1994) 417.
- [27] M. Misiak, *Nucl. Phys.* **B393** (1993) 23, Erratum **B439** (1995) 461.
- [28] T. M. Aliev, and E. Iltan, hep-ph/9803272, (1998).
- [29] P. Cho and Misiak, *Phys. Rev. D* **49** (1994) 5894.
- [30] F. M. Borzumati and C. Greub, *Phys. Rev.* **D58** (1998) 0784004
- [31] G. Buchalla, G. Isidori and S. J. Rey, *Nucl. Phys.* **B511** (1998) 594.
- [32] Z. Ligeti, I. W. Steward and M. B. Wise, *Phys. Lett.* **B420** (1998) 359.
- [33] F. Krüger and L. M. Sehgal *Phys. Lett.* **B380** (1996) 199.
- [34] A. Kagan, M. Neubert, hep-ph/9805303, 1998.
- [35] N. Cabibbo and R. Gatto, *Phys. Rev.* **D124** (1961) 1577,
 H. Burkhardt, F. Jegerlehner, G. Penso and C. Verzegnassi, *Z. Phys.* **C43** (1989) 497.
- [36] F. Krüger and L. M. Sehgal *Phys. Lett.* **B380** (1996) 199.
- [37] T. M. Aliev, and E. Iltan, *Phys. Rev.* **D58** (1998) 095014.
- [38] M. S. Alam, CLEO Collaboration, to appear in ICHEP98 Conference (1998)

- [39] D. Atwood, L. Reina and A. Soni, *Phys. Rev.* **D55** (1997) 3156.
- [40] Glenn et al., *Phys. Rev. D* **80** (1998) 2289.




Figure 1: Differential Br as a function of q^2 for fixed $\bar{\xi}_{N,bb}^D = 40 m_b$ in the region $|r_{tb}| \ll 1$, at the scale $\mu = m_b$. Here solid line corresponds to the model III with LD effects, and dashed line to the SM without LD effects.




Figure 2: The same as Fig 1, but at the region $r_{tb} \gg 1$. Dotted dashed line corresponds to the model III without LD effects.




Figure 3: The same as Fig 1, but A_{FB} versus q^2 .




Figure 4: The same as Fig 2, but A_{FB} versus q^2 .




Figure 5: The same as Fig 1, but P_L versus q^2 .




Figure 6: P_L as a function of q^2 for fixed $\bar{\xi}_{N,bb}^D = 40 m_b$ in the region $r_{tb} \gg 1$, at the scale $\mu = m_b$. Here solid line corresponds to the model III with LD effects, and dashed line to the model III without LD effects.